

Natural Remedies for High Blood Pressure

Lifestyle changes and natural remedies may help to control high blood pressure, but your doctor may also recommend medication to lower high blood pressure. It is important to work with your doctor, because [untreated high blood pressure](#) may damage organs in the body and increase the risk of heart attack, stroke, [brain hemorrhage](#), kidney disease, and [vision loss](#). See a [drawing of a hypertensive heart](#).

- **Coenzyme Q10 (CoQ10)**

There is some evidence that the supplement CoQ10 may help to reduce high blood pressure.

A 12 week double-blind, placebo-controlled trial of 83 people with systolic hypertension examined the effect of CoQ10 supplements (60 mg twice daily). After the 12 weeks, there was a mean reduction in systolic blood pressure of 17.8 mm Hg in the Coq10-treated group.

Another study conducted at the University of Western Australia looked at the effect of CoQ10 on blood pressure and glycemic control in 74 people with type 2 diabetes. Participants were randomly assigned to receive either 100mg CoQ10 twice daily, 200mg of the drug fenofibrate, both, or neither for 12 weeks.

CoQ10 significantly reduced systolic and diastolic blood pressure (mean reduction 6.1 mm Hg and 2.9 mm Hg respectively). There was also a reduction in HbA1C, a marker for long-term glycemic control. To learn more about CoQ10, read the [Coenzyme Q10 \(CoQ10\) fact sheet](#).

- **Garlic**

In a meta-analysis of seven randomized controlled trials of garlic supplements, three trials showed a significant reduction in systolic blood pressure and four in diastolic blood pressure. Researchers concluded that garlic powder supplement may be of clinical use in patients with mild high blood pressure.

Garlic supplements should only be used under the supervision of a qualified health practitioner. Garlic can thin the blood (reduce the ability of blood to clot) similar to aspirin. Garlic may interact with many drugs and supplements such as the prescription "blood-thinners" drugs such as Coumadin (warfarin) or Trental (pentoxifylline), aspirin, vitamin E, ginkgo. It is usually recommended that people taking garlic stop in the weeks before and after any type of surgery.

To learn more about garlic, go to the [articles about garlic](#).

- **Hawthorn**

The herb hawthorn is often used by traditional herbal practitioners for high blood pressure.

In a randomized controlled trial conducted by researchers in Reading, UK, 79 patients with type 2 diabetes were randomized to receive either 1200 mg of hawthorn extract a day or placebo for 16 weeks. Medication for high blood pressure was used by 71% of the patients.

At the end of the 16 weeks, patients taking the hawthorn supplement had a significant reduction in mean diastolic blood pressure (2.6 mm Hg). No herb-drug interactions were reported.

- **Fish oil**

Preliminary studies suggest that fish oil may have a modest effect on high blood pressure. Although fish oil supplements often contain both DHA (docosahexaenoic acid) and EPA (eicosapentaenoic acid), there is some evidence that DHA is the ingredient that lowers high blood pressure. Learn more about [fish oil](#).

- **Folic acid**

Folate is a B vitamin necessary for formation of red blood cells. It may help to lower high blood pressure in some people, possibly by reducing elevated homocysteine levels.

One small study of 24 cigarette smokers found that four weeks of folic acid supplementation significantly lowered blood pressure. Learn more about [folic acid](#).

- **Diet**

Changing your diet is an important part of lowering high blood pressure. The DASH (Dietary Approaches to Stop Hypertension) diet is promoted by the National Heart, Lung, and Blood Institute of the National Institutes of Health (NIH).

The [DASH diet](#) includes fruits and vegetables, low-fat dairy foods, beans and nuts. Sodium is limited to 2,400 mg per day.

Studies have found that the DASH diet can reduce high blood pressure within two weeks. These are the daily guidelines of the DASH diet:

- 7 to 8 servings of grains
- 4 to 5 servings of vegetables
- 4 to 5 servings of fruit
- 2 to 3 servings of low-fat or non-fat dairy
- 2 or less servings of meat, fish, or poultry
- 2 to 3 servings of fats and oils
- 4 to 5 servings per week of nuts, seeds, and dry beans
- Less than 5 servings a week of sweets

Serving Sizes

1/2 cup cooked rice or pasta

1 slice bread

1 cup raw vegetables or fruit

1/2 cup cooked vegetables or fruit

8 oz. of milk

1 teaspoon olive oil

3 ounces cooked meat

3 ounces tofu

A related diet called the DASH-Sodium reduces sodium to 1,500 mg a day, which is approximately equal to 2/3 teaspoon from all sources (processed and canned foods contain hidden salt).

Patients following the DASH-Sodium diet had a significant reduction in high blood pressure.

More: [The Salt Wars: Is Salt Restriction Necessary?](#)

- **Calcium, Magnesium, and Potassium**

Calcium. Calcium supplementation appears to have a modest but statistically significant reduction in systolic blood pressure (mean difference of 2.5 mm Hg), however better quality studies are needed. Learn more about [getting enough calcium in your diet](#).

Potassium. A meta-analysis of five trials indicated that potassium supplementation compared to a control resulted in a large but statistically non-significant reduction in systolic blood pressure (mean difference 11.2 mm Hg) and diastolic blood pressure (5.0 mm Hg). Learn about [potassium in the diet](#) and find out [which foods have potassium](#).

Magnesium. In 12 randomized controlled trials, participants receiving magnesium supplements did not have a significant reduction in systolic blood pressure, but they did have a statistically significant reduction in diastolic blood pressure (mean difference 2.2 mm Hg). Read more about [magnesium in the diet](#).

- **Mind-Body Interventions**

Mind-body interventions, particularly autogenic training, biofeedback, and yoga, have been found to modestly reduce high blood pressure compared with placebo.

Autogenic Training

Autogenic training is a technique used for stress reduction and relaxation. It involves a series of sessions in which people learn how to control breathing, blood pressure, heart rate, and body temperature.

People learn six exercises that each involve a certain posture (e.g. reclining in a chair), concentration without a goal, imagination, and verbal cues. Each exercise is learned by watching a teacher demonstrate it or by reading a description. It requires regular practice.

Biofeedback

Biofeedback is a technique in which people learn how to gain control over internal body processes that normally occur involuntarily, such as blood pressure, heart rate, muscle tension, and skin temperature.

Biofeedback is primarily used for high blood pressure, migraine, tension headache, chronic pain, and urinary incontinence.

Of the different types of biofeedback, thermal feedback (which measures skin temperature) and electrodermal activity feedback (which uses a probe that responds to sweat) may be more effective than direct blood pressure feedback or electromyography (EMG), which measures muscle tension. Learn more about [biofeedback](#).

Yoga

Preliminary studies have found that yoga may lower blood pressure. Go to the [yoga resources](#) page.

- **Aerobic Exercise**

Aerobic exercise is an important part of the natural approach to lower high blood pressure. A meta-analysis of 105 trials involving a total of 6805 participants found that aerobic exercise was associated with a mean reduction in systolic blood pressure of 4.6 mm Hg, with corresponding reductions in diastolic blood pressure.

People with high blood pressure should speak with their doctor first before embarking on a new exercise program. Learn about the [effect of exercise on blood pressure and the heart](#).

• **Ayurvedic Medicine**

In Ayurveda, the traditional medicine of India, high blood pressure is treated according to each person's dosha, or constitutional type.

The pitta type may have a flushed face, red eyes, headaches, light sensitivity, irritability, and nosebleeds.

The kapha type may have excess weight, water retention, high cholesterol, sluggishness.

The vata type may feel cold, have gas, bloating, or constipation, insomnia, or nervousness, worry, or anxiety.

[What's Your Ayurvedic Type?](#)
[Ayurvedic Foods for Each Type](#)
[Ayurveda Factsheet](#)

• **Traditional Chinese Medicine**

In traditional Chinese medicine, high blood pressure is often attributed to a problem with the circulation of vital energy (qi) in the body. Chinese medicine practitioners believe that depression, anger, obesity, and high intake of fatty foods are some of the causative factors.

A combination of acupuncture and herbs is often recommended. Foods thought to have medicinal properties that may help high blood pressure include water chestnut, turnip, honey, Chinese celery, hawthorn berries, and mung beans.

Emotions also play a role in our health. Learn about [emotions in traditional Chinese medicine](#).

Sources:

Ernst E. "Complementary/alternative medicine for hypertension: a mini-review." *Wien Med Wochenschr.* 155.17-18 (2005): 386-391.

Dickinson HO, Mason JM, Nicolson DJ, Campbell F, Beyer FR, Cook JV, Williams B, Ford GA. Lifestyle interventions to reduce raised blood pressure: a systematic review of randomized controlled trials. *J Hypertens.* 24.2 (2006): 215-233.

Dickinson HO, Nicolson DJ, Campbell F, Beyer FR, Mason J. "Potassium supplementation for the management of primary hypertension in adults." *Cochrane Database Syst Rev.* 19.3 (2006): CD004641.

Dickinson HO, Nicolson DJ, Campbell F, Cook JV, Beyer FR, Ford GA, Mason J. "Magnesium supplementation for the management of essential hypertension in adults." *Cochrane Database Syst Rev.* 19.3 (2006): CD004640.

Dickinson HO, Nicolson DJ, Cook JV, Campbell F, Beyer FR, Ford GA, Mason J. Calcium supplementation for the management of primary hypertension in adults. *Cochrane Database Syst Rev.* 19.2 (2006): CD004639.

Geleijnse JM, Giltay EJ, Grobbee DE, et al. "Blood pressure response to fish oil supplementation: metaregression analysis of randomized trials." *J Hypertens.* 20 (2002): 1493-1499.

He FJ, MacGregor GA. "Effect of longer-term modest salt reduction on blood pressure." *Cochrane Database Syst Rev.* 3 (2004): CD004937.

Hodgson JM, Watts GF, Playford DA, Burke V, Croft KD. "Coenzyme Q10 improves blood pressure and glycaemic control: a controlled trial in subjects with type 2 diabetes." *Eur J Clin Nutr.* 56.11 (2002): 1137-1142.

Jurgens G, Graudal NA. "Effects of low sodium diet versus high sodium diet on blood pressure, renin, aldosterone, catecholamines, cholesterols, and triglyceride." *Cochrane Database Syst Rev.* 1(2004): CD004022.

Linden W, Moseley JV. "The efficacy of behavioral treatments for hypertension." *Appl Psychophysiol Biofeedback.* 31.1 (2006): 51-63.

Mangoni AA, Sherwood RA, Swift CG, Jackson SH. "Folic acid enhances endothelial function and reduces blood pressure in smokers: a randomized controlled trial." *J Intern Med.* 252.6 (2002): 497-503.

Martinez-Mir I, Rubio E, Morales-Olivas FJ, Palop-Larrea V. "Transient ischemic attack secondary to hypertensive crisis related to Panax ginseng." *Ann Pharmacother.* 28.11 (2004): 1970.

Mori TA, Bao DQ, Burke V, et al. "Docosahexaenoic acid but not eicosapentaenoic acid lowers ambulatory blood pressure and heart rate in humans." *Hypertension.* 43 (1999): 253-260.

Silagy CA, Neil HA. "A meta-analysis of the effect of garlic on blood pressure." *J Hypertens.* 12.4 (1994): 463-468.

Walker AF, Marakis G, Simpson E, Hope JL, Robinson PA, Hassanein M, Simpson HC. "Hypotensive effects of hawthorn for patients with diabetes taking prescription drugs: a randomised controlled trial." *Br J Gen Pract.* 56. 527 (2006): 437-443.

Williams C, Kingwell BA, Burke K, McPherson J, Dart AM. Folic acid supplementation for 3 wk reduces pulse pressure and large artery stiffness independent of MTHFR genotype. *Am J Clin Nutr.* 82.1 (2005): 26-31.

Reference website: <http://altmedicine.about.com/cs/herbsvitaminsek/a/Hypertension.htm>